

# Management *Guide*

LOHMANN TIERZUCHT

## Parent Stock

LOHMANN BROWN



LOHMANN LSL

## PRODUCTS OF LOHMANN TIERZUCHT

**LOHMANN LSL-CLASSIC**

**LOHMANN BROWN-CLASSIC**

**LOHMANN LSL-LITE**

**LOHMANN BROWN-LITE**

**LOHMANN LSL-EXTRA**

**LOHMANN BROWN-EXTRA**

**LOHMANN TRADITION**

**LOHMANN SANDY**

**LOHMANN SILVER**


Increasing worldwide concentration and growing competition in the poultry industry requires efficient layers to satisfy specific market requirements.

LOHMANN TIERZUCHT offers a wide range of high quality layer strains "bred in Germany" to meet these demands.

The intensive monitoring programme of all breeding farms and hatcheries by our Veterinary Laboratory assures the highest possible health status of chicks supplied by LOHMANN TIERZUCHT.



The mainstream products are **LOHMANN LSL-CLASSIC** and **LOHMANN BROWN-CLASSIC**, well known for their efficient production of quality white and brown eggs, respectively.



**LOHMANN LSL-LITE** and **LOHMANN BROWN-LITE** are two products, designed for markets which prefer smaller eggs and measure efficiency in g feed per egg.

## PRODUCTS OF LOHMANN TIERZUCHT



For markets requiring more XL-size eggs **LOHMANN-LSL-EXTRA** and **BROWN-EXTRA** are the ideal white and brown layers.



**LOHMANN SANDY** is a white feathering layer for the production of cream coloured eggs. The layer has an outstanding feed conversion and robustness.



**LOHMANN TRADITION**, a brown egg layer with high early egg weight is being developed mainly for markets requiring even bigger egg size.



**LOHMANN SILVER** is a predominately white feathering layer for the production of uniform brown eggs with reduced egg weight. Her special advantage is the excellent feathering.

# CONTENTS



<b>4</b>	<b><i>Introduction</i></b>
<b>5</b>	<b><i>Top Performance by Systematic Selection</i></b>
<b>6</b>	<b><i>Breeding Scheme Lohmann LSL</i></b>
<b>7</b>	<b><i>Performance Data Lohmann LSL Parent Stock</i></b>
<b>8</b>	<b><i>Breeding Scheme Lohmann Brown</i></b>
<b>9</b>	<b><i>Performance Data Lohmann Brown Parent Stock</i></b>
<b>10</b>	<b><i>Housing Chicks</i></b>
10	General Recommendations
10	Cage Systems
10	Floor Systems
11	Environment
12	Brooding Temperature
12	Body Temperature of the Chicks
<b>13</b>	<b><i>Vaccination</i></b>
13	General Recommendations
13	Vaccination Methods
13	Special Recommendations
14	Example of a Vaccination Programme
<b>15</b>	<b><i>Beak Treatment</i></b>
<b>16</b>	<b><i>Nutrition</i></b>
16	Principles
19	Rearing and Growing Period
22	Laying Period
24	Feed Consumption
24	Micro-Nutritions
24	Heat Treatment

# CONTENTS



<b>26</b>	<b><i>Lighting</i></b>
26	General
26	Intermittent Lighting Programme for Day Old Chicks
27	Lighting Programme for Closed Houses
29	Lighting Programme for Open Houses
30	Light Intensity
<b>33</b>	<b><i>General Recommendations</i></b>
33	Hygiene
33	Insect and Parasite Control
33	Cleaning and Disinfection
34	Daily Control
34	Mating Ratio
34	Water Supply
35	Grit
35	Stocking Density
35	Equipment Requirements
36	Litter
36	Nests and Egg Collection
37	Hatching Egg Care
37	Hatching Egg Disinfection
<b>38</b>	<b><i>General Information</i></b>
38	Body Weight Development of Lohmann LSL Parent Stock
39	Body Weight Development of Lohmann Brown Parent Stock
40	Performance Goals of Lohmann LSL Parent Stock
42	Performance Goals of Lohmann Brown Parent Stock

# INTRODUCTION

## Introduction

LOHMANN LSL and LOHMANN BROWN parent stock are bred to produce high performance layers for profitable egg production. The source lines are carefully selected for combining ability.

Each parent flock is a significant investment. To maximise return on investment, good management practices are required. This guide contains management recommendations based on comprehensive international experience. Local conditions may require specific adaptations of management practices to achieve optimal results.

Genetic selection within the pure lines is a continuing process to assure that LOHMANN LSL and LOHMANN BROWN

parents and commercial layers meet market requirements and produce more profit than other strains.

Egg quality at the parent and commercial level depends on a combination of genetic potential and non-genetic factors (health, nutrition, light, temperature, air quality, technical environment). It pays to control the non-genetic variables in order to help the birds express their genetic potential.

We trust that our management recommendations in this guide for LOHMANN LSL and LOHMANN BROWN parent stock will help to improve your technical results from flock to flock. Keep accurate records and contact your area representative of LOHMANN TIERZUCHT for more detailed information. ■



# TOP PERFORMANCE BY SYSTEMATIC SELECTION



In recent decades advanced methods have significantly improved breeding quality. Due to the development of powerful electronic data processing systems, it has become possible to put the theory of selection systematically into practice – thus turning modern quantitative genetics into reality.

From very early on, LOHMANN TIERZUCHT used these new techniques and can therefore offer an extensive range of experience and know-how. A highly qualified team of specialists guarantees prompt utilization of the latest research results. The market's changing demands can therefore be met quickly and effectively.

Moreover, nationally and internationally, LOHMANN TIERZUCHT is ranked as first class for questions on poultry health, which is one of the decisive factors for performance and profitability.

Intensive research in our own Veterinary Laboratory, besides increasing resistance to diseases by genetic means and ensuring the strictest conditions of hygiene, is fundamental to the quality of LOHMANN TIERZUCHT products. In addition, LOHMANN TIERZUCHT also provides expert advice on all questions of feed, nutrition and technical service.


Practice profits from this extensive expertise in all aspects of poultry management. With LOHMANN TIERZUCHT products, eggs are produced in top quality and at competitive costs.

Results of performance comparisons in the field and in independent institutes are proof of this success. LOHMANN TIERZUCHT products are often the winners and are always among the few at the top, worldwide.

**LOHMANN TIERZUCHT – the right partner for progressive, successful poultry management. ■**

## BREEDING SCHEME

### LOHMANN LSL



## PERFORMANCE DATA



### LOHMANN LSL Parent Stock


<b>Egg Production</b>	Age at 50% production	21 – 22 weeks
	Peak production	26 – 30 weeks
	<b>Eggs per Hen Housed</b>	
	to 68 weeks of age	280 – 290
	to 72 weeks of age	295 – 305
<b>Settable Eggs per Hen Housed</b>	to 68 weeks of age	250 – 260
	to 72 weeks of age	270 – 280
	<b>Saleable Chicks per Hen Housed</b>	
	to 68 weeks of age	100 – 110
	to 72 weeks of age	110 – 115
<b>Hatchability</b>	Average	80 – 83 %
<b>Body Weight</b>	Female	
	at 20 weeks	1.2 – 1.4 kg (2.6 – 3.1 lbs.)
	at 72 weeks	1.6 – 1.8 kg (3.5 – 4.0 lbs.)
	Male	
	at 20 weeks	1.6 – 1.8 kg (3.5 – 4.0 lbs.)
	at 72 weeks	2.2 – 2.4 kg (4.8 – 5.3 lbs.)
<b>Feed Consumption (Male and Female)</b>	1 – 20 weeks	7.2 kg (15.8 lbs.)
	21 – 72 weeks	42 kg (92.6 lbs.)
<b>Liveability</b>	Rearing	96 – 98 %
	Laying period	90 – 96 %





## BREEDING SCHEME

### LOHMANN BROWN



## PERFORMANCE DATA



### LOHMANN BROWN Parent Stock

<b>Egg Production</b>	Age at 50 % production	22 – 23 weeks
	Peak production	26 – 30 weeks
	<b>Eggs per Hen Housed</b>	
	to 68 weeks of age	270 – 280
	to 72 weeks of age	285 – 295
<b>Settable Eggs per Hen Housed</b>	to 68 weeks of age	245 – 255
	to 72 weeks of age	260 – 270
	<b>Saleable Chicks per Hen Housed</b>	
	to 68 weeks of age	100 – 105
	to 72 weeks of age	105 – 110
<b>Hatchability</b>	Average	80 – 82 %
<b>Body Weight</b>	Female	
	at 20 weeks	1.5 – 1.7 kg (3.3 – 3.7 lbs.)
	at 72 weeks	1.8 – 2.1 kg (3.4 – 4.6 lbs.)
	Male	
<b>Feed Consumption (Male and Female)</b>	1 – 20 weeks	8.0 kg (17.6 lbs.)
	21 – 72 weeks	43 kg (94.8 lbs.)
<b>Liveability</b>	Rearing	96 – 98 %
	Laying period	90 – 96 %



# HOUSING CHICKS



## General Recommendations

- Before bringing in the chicks, check that everything is in good working order.
- Males and females should be reared together from the first day.
- Warm up the house in good time to 35–36°C (95–97°F). In summer start heating at least 24 hours and in winter at least 48 hours before the chicks arrive. When the right temperature has been achieved, supply minimum ventilation. This will avoid temperature differences within the house.
- Maintain the recommended temperatures (35–36°C/95–97°F) during the first 48–72 hours.
- Relative humidity should be at least 60%.
- The right height of the drinkers must be adjusted to allow the chicks to drink water without difficulty.
- Reduce the water pressure of the nipples in order to enable the chicks to find water easily.
- Keep drinking water temperature between 20–25°C (68–77°F) by temporarily flushing the nipple drinker lines or renewing the water in the chick founts.

Follow the recommended lighting programmes (refer to page 26).

## Cage Systems

- Adjust the cage floors and feeding grids according to the manufacturer's instructions.
- Place sheets of paper on the cage floor for the first days and distribute a bit of feed on this paper. The papers must be removed by day 7.
- Unload all chick boxes and distribute them in the house. Remove all lids and place them on the top of the boxes.
- Quickly place the chicks near feeders and drinkers. Distribute the chicks evenly among the cages starting at the far end of the house.
- Trigger nipples/water cups to encourage birds to drink.

## Floor Systems

- Before arrival of the chicks, litter should be spread only after heating the house, when the floor has reached the optimum temperature. Softwood shavings or straw make suitable litter.
- After arrival, place chicks under brooders as soon as possible.
- Measure the brooder temperature by placing the thermometer 8 cm (3.15 in) inside the outer edge of the brooder and 8 cm (3.15 in) above the litter.

# HOUSING CHICKS



- Dip the beaks of a few chicks into water and trigger nipple or water cups to help them start drinking. When drinking water has been found by all chicks (this will take approx. 2–3 hours), they will start to eat.
- Supply the chicks with additional feeding bowls to ensure a better feed intake in the first few days.
- Chicks should be fully feathered before brooding equipment has to be removed.

## Environment

Environmental conditions have an effect on the wellbeing and performance of the birds. Important environmental factors are temperature, humidity and level of toxic gases in the air. The optimal temperature depends on the age of the birds. The following table is a guide to the recommended temperature at bird level.

After a few hours, check whether the chicks have settled down well. The chicks' behaviour is the best indicator of their well being:

- If the chicks are evenly spread out and moving freely, temperature and ventilation are all right.
- If the chicks are crowding together or avoiding certain areas within the

house, temperature is too low or there is a drought.

- If the chicks are laying on the floor with their wings spread out and gasping for air, temperature is too high.

**At first signs that the chicks are not feeling well determine the reason, correct the situation and check more frequently.**

**Table 1: Desired Temperatures at Bird Level Dependent on Age**

Age	Cage Rearing		Floor Rearing	
	°C	°F	°C	°F
Day 1–2 *	35	95	36	97
Day 3–4	33	91	34	93
Day 5–7	31	88	32	90
Week 2	28	82	29	84
Week 3	26	79	27	81
Week 4	22	72	24	75
From Week 5	18–20	64–68	18–20	64–68

*\* Body temperatures of 40–41 °C (104–106 °F) are the optimum for the chicks.*

## HOUSING CHICKS



The air quality should meet the following minimum requirements:

**Table 2: Minimum Air Quality Requirements**

O <sub>2</sub>	over	20 %
CO <sub>2</sub>	under	0.3 %
CO	under	40 ppm
NH <sub>3</sub>	under	20 ppm
H <sub>2</sub> S	under	5 ppm

### Brooding Temperature

Always reduce temperature gradually, and avoid sudden changes.

If the ventilation system is used to regulate temperature, take care that the necessary fresh air is supplied. The relative humidity inside the house should be 60 – 70 %.

### Body Temperature of the Chicks

There are findings which confirm that the temperature of chicks is between 40.0 (104°F) and 41.0°C (105.8°F) after the moment of full homeothermy. This information can be parallelly used with the behaviour of the housed chicks to

adjust house temperatures in an optimal way. Use modern ear thermometers, known from human medicine, as these are useful devices to measure the body temperature of day old chicks.



Make sure that you collect samples of chicks in different parts of the house and control the rectal temperature of the latter. Proceed in a way like you normally would do when weighing chicks/pullets and check for uniformity. Obtain samples from chicks distributed throughout the house in order to have reliable readings. Collect the information, calculate the average and adjust the house temperatures accordingly to achieve optimal chick temperatures.

If the actual barn temperature, humidity or uniformity of air distribution are significantly below the recommended levels, chick growth maybe adversely affected due to chilling. ■

## VACCINATION



### General Recommendations

Vaccination is an important way of preventing diseases. Different regional epidemic situations require suitably adapted vaccination programmes. Therefore, please be guided, by the advice of your local veterinarian and poultry health service. Only healthy flocks should be vaccinated. Check the expiration date of the vaccine. The vaccine must not be used after this date. Keep records of all vaccinations and vaccine serial numbers.

### Vaccination Methods

**Individual Vaccinations** such as injections and eye-drops are very effective and generally well tolerated but also very labour intensive.

**Drinking Water Vaccinations** are not labour intensive but must be carried out with the greatest care to be effective. The water used for preparing the vaccine solution must not contain any disinfectants. During the growing period, the birds should be without water for approximately 2 hours prior to vaccination. During hot weather reduce this time accordingly. The amount of vaccine solution should be calculated for complete consumption within 2–4 hours. When vaccinating with live vaccines, add 2g of skim milk powder per litre of water or canned milk in order to protect the virus titre, if no water stabilisator is available.

**Spray Vaccinations** are not labour intensive and are highly effective, but may occasionally have side effects. For chicks up to the age of 3 weeks apply only coarse spray. Use distilled water for vaccination.

### Special Recommendations

**Marek Re-Vaccinations** have proved to be successful after long transportation and in areas with high infection risk. Consult your veterinarian and the Lohmann Veterinary Laboratory for further information.

**Mycoplasmosis Vaccinations** are only advisable if the farm cannot be kept free of mycoplasmosis. Infections with virulent mycoplasma species during the production period lead to performance depression. The best performance is achieved by flocks which are kept free of mycoplasmosis and are not vaccinated.

**Vaccination against Coccidiosis** is the most reliable method in the floor rearing to develop immunity against this disease. Never use coccidiostats in the feed when pullets are vaccinated.

**Applying Vitamins** in the first two to three days after vaccination can help to reduce stress and prevent undesired reactions. To what extent depends on the specific situation on each farm.



## VACCINATION



Table 3: Example of a Vaccination Programme

Disease	Occurrence		Application Methods	Remarks
	World-wide	Locally		
Marek	●		SC – IM	Day 1 – Hatchery
Newcastle*	●		DW – SP – SC – IM	Number of vaccinations according to disease pressure
Gumboro (IBD)*	●		DW	2 live vaccinations recommended
Infectious Bronchitis*	●		DW – SP – SC – IM	Number of vaccinations according to disease pressure
AE	●		DW – SC – WW	Vaccination between 8 – 14 weeks
CAV	●		DW – SC – IM	Vaccination between 8 – 14 weeks
Mycoplasmosis		●	SP – ED – SC – IM	Vaccination before transfer
Fowl Pox		●	WW	Vaccination before transfer
Pasteurellosis		●	SC	2 vaccinations between 8 – 14 weeks
Infectious Coryza		●	SC	2 vaccinations between 8 – 14 weeks
Salmonella		●	DW – SP – IM	Vaccination before transfer
ILT		●	DW – ED	2 vaccinations between 6 – 14 weeks
EDS		●	SC – IM	Vaccination before transfer
E. Coli		●	SC – IM	2 vaccinations between 6 – 16 weeks

■ DW: Drinking Water ■ SP: Spray ■ ED: Eye Drop ■ WW: Wing Web ■ IM: Intramuscular Injection  
 ■ SC: Subcutaneous Injection ■ Vaccination against Coccidiosis is optional for floor rearing systems

\*An implementation of early live vaccination for Newcastle Disease (ND) and Infectious Bronchitis (IB) is of high value in order to induce local protection in the respiratory system of the chicks (priming effect). The right choice of vaccine is crucial. Never vaccinate very young birds with high-virulence live vaccine. Revaccination with live ND and/or IB every 6–8 weeks during production period is beneficial in order to improve the local immunity. The use of inactivated ND/IB/IBD vaccine before onset of lay is recommended.

## BEAK TREATMENT



Beak treatment is not necessary under optimal conditions. In practice, it is widely used in environmental controlled and light-tight facilities, as an efficient precaution against cannibalism and feather pecking. Such behaviour may develop at any age as a result of excessive light intensity, unbalanced feed, undersupply of nutrients, poor ventilation, overstocking or boredom.

Especially in floor management and/or open houses with uncontrollable light intensity, we recommend beak treatment subject to local animal welfare regulations.

A very gentle and highly recommended method of beak treatment is the infrared treatment of the upper and lower beak by means of a special technique, performed shortly after chicks hatch. This procedure can already be done in the hatchery under very hygienic conditions by specially trained personnel. Another method of beak treatment is to treat the beaks with a hot blade.

### Observe the following precautions for a conventional beak treatment:

- Treat only healthy, unstressed birds, at the age of 7 – 10 days.
- Allow only experienced personnel to do the work.
- Work slowly and carefully.

- Use only equipment and blades in perfect working order; adjust the blade temperature so that cauterisation is guaranteed and the beak is not damaged.
- Adjust temperature and duration of the treatment according to the chicks beak size, strength and quality.
- Do not feed for 12 hours before treating.
- Offer free feeding immediately after treating.
- Increase the level of feed in the troughs.
- Increase the temperature in the house for a few days after treating.
- For 3–5 days after beak treating provide an extra hour of light and supply feed in the late evening or at night.
- Giving vitamins via the drinking water can also help to alleviate stress.

- Under optimal housing conditions, males should generally not be beak treated. Nevertheless, if a beak treatment for males is necessary, just treat them very gently. ■

# NUTRITION



## Principles

Correct nutrition is a prerequisite to utilize the high genetic performance potential of LOHMANN LSL and LOHMANN BROWN Parent Stock. Best control on nutrition is achieved by feeding a ready mixed compound feed. The recommended feeding programme concentrates on the essential nutrients and is

designed to cover the requirements for top performance in rearing and production.

The base for correct nutrition is correct feedstuff evaluation. The following table gives information on the sources used for the recommendations in this programme and on analytical methods for the determination of specific nutrients.

Nutrient	Source of Information	Analytical Method
Energy	Results of European Research Institutes	N-corrected Metabolizable Energy
Crude Protein	Results of European Research Institutes	Method according to Kjeldahl
Amino Acids	Results of LOHMANN TIERZUCHT and Amino Acid Producers	Amino Acid Analyser
Digestible Amino Acids	Digestion Coefficients based on Dutch trials	
Calcium Sodium	Results of LOHMANN TIERZUCHT	Flame - Photometry
Total Phosphorus	Results of LOHMANN TIERZUCHT	Ultraviolet-Spectroscopy
Available Phosphorus	Results of French Availability Studies	
Chlorine	Results of LOHMANN TIERZUCHT	Titration Method acc. to Mohr
Linoleic Acid	Results of LOHMANN TIERZUCHT	Gas-Chromatography Analysis

# NUTRITION



**Table 4: LOHMANN LSL Parent Stock Females**  
Body Weight Development, Feed Consumption and Water Consumption with Standard Lighting Programme

Age in Weeks	Body Weight (g)			kJ Bird/Day	Feed Consumption		Water Consumption	
	Average	Range			g/Bird/Day	Cumulative (g)	ml/Bird/Day	Cumulative (ml)
1	65	62	68	126	11	74	18	126
2	130	123	137	204	17	193	33	357
3	190	180	200	276	23	354	44	665
4	250	237	263	319	28	550	53	1036
5	320	303	337	365	32	774	61	1463
6	400	379	421	410	36	1026	70	1673
7	485	460	510	456	40	1306	79	2226
8	570	540	600	502	44	1614	88	2842
9	650	616	684	547	48	1950	95	3507
10	730	692	768	593	52	2314	100	4207
11	810	767	853	638	56	2706	105	4942
12	880	834	926	684	60	3126	110	5712
13	945	895	995	730	64	3574	116	6524
14	1005	952	1058	764	67	4043	121	7371
15	1065	1009	1121	798	70	4533	126	8253
16	1120	1061	1179	832	73	5044	131	9170
17	1170	1109	1231	878	77	5583	137	10129
18	1220	1156	1284	935	82	6157	142	11123
19	1270	1203	1337	992	87	6766	147	12152
20	1320*	1251	1389	1049	92	7410	152	13216

\* Depends upon sexual maturity - 100 – 200 g higher for hens in production.

■ Variation is possible due to differences in feed consumption and environments.



**Table 5: LOHMANN BROWN Parent Stock Females**  
Body Weight Development, Feed Consumption and  
Water Consumption with Standard Lighting Programme

Age in Weeks	Body Weight (g)			kJ Bird/Day	Feed Consumption		Water Consumption	
	Average	Range			g/Bird/Day	Cumulative (g)	ml/Bird/Day	Cumulative (ml)
1	65	62	68	120	10	70	18	126
2	130	124	137	228	19	203	33	357
3	180	171	189	300	25	378	44	665
4	250	238	263	342	30	588	53	1036
5	320	304	336	399	35	833	61	1463
6	410	390	431	456	40	1113	70	1673
7	500	475	525	513	45	1428	79	2226
8	590	561	620	570	50	1778	88	2842
9	680	646	714	616	54	2156	95	3507
10	770	732	809	650	57	2555	100	4207
11	860	817	903	684	60	2975	105	4942
12	950	903	998	718	63	3416	110	5712
13	1030	979	1082	752	66	3878	116	6524
14	1110	1055	1166	787	69	4361	121	7371
15	1190	1131	1250	821	72	4865	126	8253
16	1270	1207	1334	855	75	5390	131	9170
17	1350	1283	1418	889	78	5936	137	10129
18	1440	1368	1512	923	81	6503	142	11123
19	1530	1454	1607	958	84	7091	147	12152
20	1600*	1520	1680	992	87	7700	152	13216

\* Depends upon sexual maturity - 100 – 200 g higher for hens in production.

■ Variation is possible due to differences in feed consumption and environments.



## Rearing and Growing Period

The recommended feed schedule for the rearing period of LOHMANN LSL and LOHMANN BROWN Parent Stock is based on four diets.

The Starter is a diet with a high nutrient density, based on a feed formulation including raw materials of excellent quality and digestibility. This feed is supposed to be used until chicks have reached body weight targets in the first three weeks.

It is followed by a traditional Grower feed based on an energy level of 11.4 MJ/kg (2720 Kcal/kg). This feed is supposed to be fed till chicks have reached a body weight targets at 8 weeks of age.

A Developer feed should be fed after 8 weeks. A low nutrient density, a good feed structure and crude fibre content up to 5 – 6 % in this special feed should be used to develop eating capacities.

The use of a Pre-Layer feed for LOHMANN LSL and LOHMANN BROWN Parent Stock has several advantages:

- The Pre-Layer feed gives a better uniformity due to the higher protein and amino acid content in the critical period of sexual maturity. Individual males and females with weight below standard are given the chance to compensate.

- The Pre-Layer feed has a higher calcium content than the Developer and improves the shell quality of early maturing hens at a later age.
- The Pre-Layer feed also smoothens the transition from low to high Calcium amount in the feed.

**Please consider the following recommendations while using Pre-Layer feed:**

- Start using Pre-Layer feed dependent on to the birds' sexual maturity, age and their standard body weights.
- Use Pre-Layer feed for about 10 days with a maximum of 1 kg per bird.
- The wrong way to use Pre-Layer feed is either to start using it too early and/or use it too long.

## NUTRITION



Table 6: Recommended Nutrient Levels (Rearing)

Diet type*		Starter	Grower	Developer	Pre-Layer
Nutrient		1 – 3 weeks	4 – 8 weeks	9 – 17/18 weeks	18/19 weeks – 5 % prod.
Metabol. Energy	kcal	2900	2720 – 2800	2720 – 2800	2720 – 2800
	MJ	12.00	11.40 – 11.70	11.40 – 11.70	11.40 – 11.70
Crude Protein	%	20.00	18.50	15.00	17.50
Methionine	%	0.48	0.40	0.34	0.36
Dig. Methionine	%	0.39	0.33	0.28	0.29
Meth./Cysteine	%	0.83	0.70	0.60	0.68
Dig. M/C	%	0.68	0.57	0.50	0.56
Lysine	%	1.20	1.00	0.68	0.85
Dig. Lysine	%	0.98	0.82	0.55	0.70
Valine	%	0.89	0.75	0.53	0.64
Dig. Valine	%	0.76	0.64	0.46	0.55
Tryptophan	%	0.23	0.21	0.17	0.20
Dig. Tryptophan	%	0.19	0.17	0.14	0.16
Threonine	%	0.80	0.70	0.55	0.60
Dig. Threonine	%	0.65	0.57	0.44	0.49
Isoleucine	%	0.83	0.75	0.60	0.74
Dig. Isoleucine	%	0.68	0.62	0.50	0.61
Calcium	%	1.05	1.00	0.90	2.00
Phosphorus, total	%	0.75	0.70	0.58	0.65
Phosphorus, avail.	%	0.48	0.45	0.37	0.45
Sodium	%	0.18	0.17	0.16	0.16
Chloride	%	0.20	0.19	0.16	0.16
Linoleic Acid	%	2.00	1.40	1.00	1.00

\* The basis for switching between diet types is the hens' body weight development. The correct time for changing the diet is determined not by age, but by body weight. Chicks and pullets should therefore be weighed at regular intervals.

## NUTRITION



Table 7: Recommended Particle-Size Distribution for Chick Starter, Grower, Developer and Layer Feed (MASH)

Sieve Size	Passing Part	Sieve Size Interval	Part of Interval
0.5 mm	19 %	0 – 0.5 mm	19 %
1.0 mm	40 %	0.51 – 1.0 mm	21 %
1.5 mm	75 %	1.01 – 1.5 mm	35 %
2.0 mm	90 %	1.51 – 2.0 mm	15 %
2.5 mm	100 %	> 2 mm	10 % *
			100 %

\* Individual Particles not bigger than  
 • 3 mm in chick starter diets  
 • 5 mm in grower, developer and layer





## Laying Period

For maximum hatching egg production and optimum hatchability a phase feeding programme is recommended. Since

the requirements for specific nutrients like essential amino acids, calcium, available phosphorus and linoleic acid are changing with age, a programme with at least 2 phases is recommended.

**Table 8: Recommended Nutrient Levels per kg of Feed for different daily Feed Consumption, Phase 1 ( 20 to approx. 50 weeks)**

Nutrient		Requirement g/Hen/Day	Daily Feed Consumption			
			105 g	110 g	115 g	120 g
Metabol. Energy	kcal	2720 – 2800	2720 – 2800	2720 – 2800	2720 – 2800	2720 – 2800
	MJ	11.4 – 11.7	11.4 – 11.7	11.4 – 11.7	11.4 – 11.7	11.4 – 11.7
Protein	%	19.20	18.29	17.45	16.70	16.00
Calcium	%	4.10	3.90	3.73	3.57	3.42
Phosphorus*	%	0.63	0.60	0.57	0.55	0.52
Av. Phosphorus	%	0.44	0.42	0.40	0.38	0.37
Sodium	%	0.17	0.16	0.15	0.15	0.14
Chloride	%	0.17	0.16	0.15	0.15	0.14
Lysine	%	0.87	0.82	0.79	0.75	0.72
Dig. Lysine	%	0.71	0.68	0.65	0.62	0.59
Methionine	%	0.44	0.42	0.40	0.38	0.37
Dig. Methionine	%	0.36	0.34	0.33	0.31	0.30
Meth./Cyst.	%	0.80	0.76	0.72	0.69	0.66
Dig. M/C	%	0.65	0.62	0.59	0.57	0.54
Arginine	%	0.92	0.87	0.83	0.80	0.76
Dig. Arginine	%	0.75	0.72	0.68	0.65	0.63
Valine	%	0.74	0.71	0.68	0.65	0.62
Dig. Valine	%	0.63	0.60	0.57	0.55	0.53
Tryptophan	%	0.18	0.17	0.17	0.16	0.15
Dig. Tryptophan	%	0.15	0.14	0.14	0.13	0.12
Threonine	%	0.61	0.58	0.55	0.53	0.51
Dig. Threonine	%	0.50	0.47	0.45	0.43	0.41
Isoleucine	%	0.69	0.66	0.63	0.60	0.58
Dig. Isoleucine	%	0.57	0.54	0.52	0.49	0.47
Linoleic Acid	%	2.00	2.00	1.82	1.74	1.67

\*without Phytase



**Table 9: Recommended Nutrient Levels per kg of Feed for different daily Feed Consumption, Phase 2 ( after approx. 50 weeks of age)**

Nutrient		Requirement g/Hen/Day	Daily Feed Consumption			
			105 g	110 g	115 g	120 g
Metabol. Energy	kcal	2720 – 2800	2720 – 2800	2720 – 2800	2720 – 2800	2720 – 2800
	MJ	11.4 – 11.7	11.4 – 11.7	11.4 – 11.7	11.4 – 11.7	11.4 – 11.7
Protein	%	18.40	17.52	16.73	16.00	15.33
Calcium	%	4.30	4.10	3.91	3.74	3.58
Phosphorus*	%	0.54	0.52	0.49	0.47	0.45
Av. Phosphorus	%	0.38	0.36	0.35	0.33	0.32
Sodium	%	0.17	0.16	0.15	0.15	0.14
Chloride	%	0.17	0.16	0.15	0.15	0.14
Lysine	%	0.85	0.81	0.78	0.74	0.71
Dig. Lysine	%	0.70	0.67	0.64	0.61	0.58
Methionine	%	0.44	0.41	0.40	0.38	0.36
Dig. Methionine	%	0.36	0.34	0.32	0.31	0.30
Meth./Cyst.	%	0.79	0.75	0.71	0.68	0.65
Dig. M/C	%	0.64	0.61	0.59	0.56	0.54
Arginine	%	0.90	0.86	0.82	0.79	0.75
Dig. Arginine	%	0.74	0.71	0.67	0.65	0.62
Valine	%	0.73	0.70	0.67	0.64	0.61
Dig. Valine	%	0.62	0.59	0.57	0.54	0.52
Tryptophan	%	0.18	0.17	0.16	0.16	0.15
Dig. Tryptophan	%	0.15	0.14	0.13	0.13	0.12
Threonine	%	0.60	0.57	0.54	0.52	0.50
Dig. Threonine	%	0.49	0.47	0.45	0.43	0.41
Isoleucine	%	0.68	0.65	0.62	0.59	0.57
Dig. Isoleucine	%	0.56	0.53	0.51	0.49	0.47
Linoleic Acid	%	1.60	1.60	1.45	1.39	1.33

\*without Phytase

In the tables with recommended nutrient levels per kg of LOHMANN LSL and LOHMANN BROWN Parent Stock feed different daily feed consumption is taken into consideration.

An average daily feed consumption of 115 g can be expected with a feed containing 11.4 MJ = 2720 kcal metabolizable energy per kg at an in-house temperature of 22 °C and good feather quality.





## Feed Consumption

The level of feed intake in the production period is mainly affected by:

- Body weight
- Performance
- House temperature: Low temperature increases the maintenance requirement for energy.
- Condition of feathering: Poor feathering condition due to management mistakes or malnutrition increases the maintenance requirement for energy.
- Feed texture: Coarse texture increases while fine texture decreases feed intake.
- Energy level: The higher the energy level of the feed, the lower the feed intake and vice versa.
- Nutrient imbalances: The hen will try to compensate for any nutrient deficits by increasing feed consumption especially in the latest stages of production.

## Micro-Nutrients

The supplementation of parent stock feed with micro-nutrients like essential

vitamins, trace elements and substances like antioxidants, coccidiostats and organic acids are essential for maximum hatching egg production and hatchability. By adding these micronutrients in suitable quantities, varying contents in the raw materials are compensated and the correct supply to the parent stock is safeguarded.

Remark: Vitamin C is synthesized by poultry normally. This vitamin is not considered as essential, but in some circumstances, like heat stress or hot climate it may be important/beneficial to add 100 – 200 mg/kg complete feed during production period.

## Heat Treatment

As a preventive measure against salmonella and other bacteria/viruses a heat treatment of the feed is sometimes applied. Depending upon the treatment technique the efficiency of specific micro-nutrients can be damaged. These micro-nutrients must therefore be added in higher quantities. According to our experience mainly Vitamin A and K<sub>3</sub> are affected. For further details contact LOHMANN TIERZUCHT.



Table 10: Recommended Micro-Nutrient Specification

Supplements per kg Feed		Starter/Grower	Developer	Pre-Layer/Phase I + II
Vitamin A	I.U.	12000	12000	15000
Vitamin D <sub>3</sub>	I.U.	2500	2500	3000
Vitamin E	mg	20 – 30**	20 – 30**	50 – 100**
Vitamin K <sub>3</sub>	mg	3***	3***	5***
Vitamin B <sub>1</sub>	mg	2	2	4
Vitamin B <sub>2</sub>	mg	8	6	10
Vitamin B <sub>6</sub>	mg	4	4	6
Vitamin B <sub>12</sub>	mcg	20	20	30
Pantothenic Acid	mg	10	10	20
Nicotinic Acid	mg	30	30	50
Folic Acid	mg	1	1	2
Biotin	mcg	100	100	200
Choline	mg	300	300	400
Antioxidant	mg	100–150**	100–150**	100–150**
Coccidiostat		as required	as required	–
Manganese*	mg	100	100	100
Zinc*	mg	60	60	60
Iron	mg	40	40	40
Copper*	mg	5	5	10
Iodine	mg	1	1	1
Selenium*	mg	0.3	0.3	0.3

\* So called "organic sources" should be considered with higher bioavailability.

\*\* according to fat addition \*\*\* double in case of heat treated feed

Table 11: Continuous Supply of Fine and Coarse Limestone (Recommended Relation in Feed)

Feed type	Fine Limestone 0 – 0.5 mm	Coarse Limestone* 1.5 – 3.5 mm
Layer Phase 1	30 %	70 %
Layer Phase 2	25 %	75 %

\* can be partly replaced by oyster shells

# LIGHTING



## General

The lighting programme controls onset of lay and affects the performance during the production period. So, within certain limits, performance can be adapted to farm specific requirements by adjusting the lighting scheme. It is easiest to follow the lighting programme in closed houses. In this case the hours of light and light intensity can be adjusted to changing needs.


## Intermittent Lighting Programme for Day Old Chicks

When the day old chicks arrive on the farm, they have already been intensively handled in the hatchery and often have a long transport to their final destination. Common practice is to give them 24 hours of light to help them recover in the first 2 or 3 days after arrival and to provide them enough time to eat and drink. In practice, it can be observed that after arrival and housing, some chicks continue to sleep whereas others start to look for feed and water. The activity of the flock will always be irregular. Especially in this phase, poultry men have difficulties interpreting the chicks' behaviour and their condition. There is a practically proven principal in splitting the day into phases of resting and activity using a special designed intermittent lighting programme. The aim is to synchronise the chicks' activities. The farmer gets a better impression

of the flock's condition and the birds are encouraged by the group's behaviour to search for water and feed.

Therefore, LOHMANN TIERZUCHT advises to give chicks a rest after they arrive at the rearing farm and then start with four hours of light followed by two hours of darkness.

## Lighting Programme after Arrival



This programme can be used for up to 7 or 10 days after arrival, then switch to the regular step down lighting programme. The use of the following lighting programme brings about the following advantages:

- The chicks will rest and/or sleep at the same time. This means that the behaviour of the chicks will be synchronised.
- Weak chicks will be stimulated by stronger ones to move as well as to eat and drink.
- The behaviour of the flock is more uniform and the monitoring of the birds is made easier.
- Mortality will decrease.

# LIGHTING



## Lighting Programme for Closed Houses

In closed houses the basic principles of lighting programmes for layers are:

- **Never increase hours of light during the growing period.**
- **Never decrease hours of light during the production period.**

To what degree lighting hours are reduced during the growing period, and the time when stimulation is started by increasing the lighting hours, are means by which performance can be adapted to farm specific requirements.

Table 12: Lighting Programme for Windowless Houses

LOHMANN LSL Parent Stock

Age (Weeks)	Hours of Light (Standard)	Light Intensity (Lux*)
Day 1 – 2 **	24	20 – 40
Day 3 – 6 **	20	20 – 30
2	18	10 – 20
3	17	5 – 10
4	16	5 – 10
5	16	5 – 10
6	16	5 – 10
7	16	5 – 10
8	16	5 – 10
9	16	5 – 10
10	15	5 – 10
11	14	5 – 10
12	13	5 – 10

Age (Weeks)	Hours of Light (Standard)	Light Intensity (Lux*)
13	12	5 – 10
14	11	5 – 10
15	10	5 – 10
16	9	5 – 10
17	9	5 – 10
18	9	5 – 10
19	9	5 – 10
20	11	20 – 30
21	12	20 – 30
22	13	20 – 30
23	14***	20 – 30
24	14***	20 – 30
25****	14***	20 – 30

\* Lux = Lumen/m<sup>2</sup>

\*\* or run an Intermittent Lighting Programme

\*\*\* According to recent research results, 14 hours of light are sufficient during production provided the house is light tight.

\*\*\*\* until the end of production

## LIGHTING



**Table 13: Lighting Programme for Windowless Houses**

LOHMANN BROWN Parent Stock

Age (Weeks)	Hours of Light (Standard)	Light Intensity (Lux*)	Age (Weeks)	Hours of Light (Standard)	Light Intensity (Lux*)
Day 1–2 **	24	20 – 40	13	9	5 – 10
Day 3–6 **	20	20 – 30	14	9	5 – 10
2	16	10 – 20	15	9	5 – 10
3	12	5 – 10	16	9	5 – 10
4	9	5 – 10	17	9	5 – 10
5	9	5 – 10	18	11	5 – 10
6	9	5 – 10	19	12	5 – 10
7	9	5 – 10	20	13	20 – 30
8	9	5 – 10	21	14***	20 – 30
9	9	5 – 10	22	14.5***	20 – 30
10	9	5 – 10	23	15***	20 – 30
11	9	5 – 10	24	15.5***	20 – 30
12	9	5 – 10	25 ****	16***	20 – 30

\* Lux = Lumen/m<sup>2</sup>

\*\* or run an Intermittent Lighting Programme

\*\*\* According to recent research results, 14 hours of light are sufficient during production provided the house is light tight.

\*\*\*\* until the end of production

## LIGHTING



### Lighting Programme for Open Houses

The principle for windowless houses **“Do not increase the hours of light during rearing period and do not reduce hours of light during production period”** also applies to “open” housing.

The effect of the natural daylight must be considered when designing lighting programmes, if natural light enters the building throughout the day.

For example in Central Europe the natural day length increases in the course of the calendar year to about 17 hours until late June and then shortens to about 8 hours until late December.

If flocks are moved to an open production house with windows that cannot be darkened, the lighting programme must be adjusted to the natural day length at the time of rehousing.

We distinguish between two variants:

1. Production starts as the natural day length decreases.
2. Production starts as the natural day length increases.

In both variations the lighting programme at 18 weeks of age should be set to a lighting period of at least 10 hours, taking the natural day length into account, and to be increased by 1 hour every week to 16 hours until

25 weeks of age. Never switch on the artificial light before 04.00 o'clock in the morning (CE time).

During the spring months the lighting programme is affected by the increase of natural day length and gradually extends to about 17 hours. When the natural day length begins to decrease in Central Europe from July, the 17-hour light period should be kept constantly until the end of the production period.

This example can be accomplished in Central Europe very simply as follows:

- 04.00\* o'clock in the morning: lights on – dimmer switch off at  $\geq 50$ –60 Lux.
- Dimmer switch on at  $\leq 50$ –60 Lux – 21.00\* o'clock in the evening lights off.

\* Central European time

These times should be varied depending on the condition of the flocks, the start of lay (production, egg size) and the facilities in the building.

If for operational reasons a different diurnal rhythm from the one described above is applied, it should not differ too much from the dawn/dusk times stated above, having regard to the diurnal rhythm of the hens.

As already mentioned, the lighting programme described here is just an example adjusted to Central European time.

## LIGHTING



If the birds are driven indoors before the end of the natural day and if the building can be darkened completely, the lighting programme for windowless laying houses should be applied.

The times for darkening the room or opening the windows are determined by the lighting programme. It is important to follow the correct sequence:


- In the evening close the windows first, then switch off the light;
- In the morning switch on the light first, then open the windows.

Contact your LOHMANN TIERZUCHT specialists for specific lighting programmes adjusted to your location, conditions and requirements.

### Light Intensity

It is very important to bear in mind that birds in an open house are exposed to very high light intensity in the natural daylight. Therefore it is crucial to prepare the pullets in the rearing for such high light intensities to avoid behavioural disorders. High frequency light sources with a bright light spectrum are recommended.

An example of a lighting programme for LOHMANN BROWN CLASSIC Parent Stock adjusted to location, condition and requirements by LOHMANN TIERZUCHT lighting programme tool



## LIGHTING



Table 14: Natural Daylight at different Latitudes

Week of Year		Approximate Duration of Natural Daylight in hours							
Hemisphere		Latitude°							
North	South	25–30	30–35	35–40	40–45	45–50	50–55	55–60	60–65
1	27	10.30	10.10	9.40	9.10	8.30	8.00	6.40	5.00
2	28	10.30	10.10	9.40	9.20	8.40	8.10	6.50	5.40
3	29	10.40	10.20	10.00	9.30	8.50	8.20	7.20	6.00
4	30	10.40	10.30	10.10	9.40	9.10	8.40	7.50	6.40
5	31	10.50	10.40	10.20	10.00	9.30	9.00	8.20	7.20
6	32	11.00	10.50	10.30	10.10	10.00	9.30	9.00	8.10
7	33	11.10	11.00	10.40	10.30	10.20	10.00	9.20	9.00
8	34	11.20	11.10	11.00	10.50	10.40	10.20	9.50	9.30
9	35	11.30	11.30	11.20	11.10	11.00	10.50	10.40	10.10
10	36	11.40	11.40	11.40	11.30	11.30	11.20	11.20	11.00
11	37	11.50	11.50	12.00	11.50	11.50	11.50	12.00	11.40
12	38	12.10	12.10	12.10	12.10	12.10	12.20	12.20	12.20
13	39	12.20	12.20	12.30	12.30	12.40	12.40	13.00	13.00
14	40	12.30	12.40	12.50	12.50	13.00	13.10	13.30	13.50
15	41	12.40	13.00	13.00	13.20	13.20	13.40	14.10	14.40
16	42	12.50	13.10	13.20	13.30	13.40	14.20	14.50	15.20
17	43	13.00	13.20	13.30	13.50	14.00	14.30	15.10	16.00
18	44	13.10	13.30	13.50	14.00	14.30	15.00	15.40	16.50
19	45	13.20	13.40	14.00	14.20	14.50	15.20	16.20	17.30
20	46	13.30	13.50	14.20	14.40	15.10	15.50	16.50	18.10
21	47	13.40	14.00	14.30	15.00	15.30	16.10	17.20	18.50
22	48	13.40	14.10	14.40	15.10	15.30	16.20	17.40	19.20
23	49	13.50	14.10	14.40	15.10	15.40	16.30	18.00	19.50
24	50	13.50	14.20	14.40	15.20	15.40	16.40	18.10	20.10
25	51	13.50	14.20	14.40	15.20	15.50	16.40	18.10	20.20
26	52	13.50	14.20	14.40	15.20	16.00	16.40	18.10	20.10



Table 15: Natural Daylight at different Latitudes

Week of Year		Approximate Duration of Natural Daylight in hours							
Hemisphere		Latitude°							
North	South	25–30	30–35	35–40	40–45	45–50	50–55	55–60	60–65
27	1	13.50	14.20	14.40	15.10	15.50	16.30	18.00	19.50
28	2	13.50	14.20	14.40	15.10	15.50	16.20	17.40	19.20
29	3	13.40	14.20	14.30	15.10	15.30	16.10	17.20	18.40
30	4	13.30	14.00	14.20	14.40	15.20	15.50	16.50	18.10
31	5	13.30	13.50	14.10	14.30	14.50	15.30	16.20	17.30
32	6	13.20	13.30	13.50	14.10	14.30	15.00	15.50	16.50
33	7	13.10	13.20	13.40	13.50	14.10	14.30	15.10	16.00
34	8	13.00	13.10	13.20	13.30	13.50	14.00	14.50	15.20
35	9	12.50	13.00	13.10	13.20	13.30	13.40	14.10	14.50
36	10	12.40	12.40	12.50	13.00	13.10	13.20	13.40	14.00
37	11	12.30	12.30	12.30	12.40	12.40	12.50	13.00	13.10
38	12	12.30	12.10	12.10	12.10	12.10	12.20	13.20	12.20
39	13	12.10	12.00	12.00	11.50	11.50	12.00	11.50	11.50
40	14	11.50	11.50	11.40	11.30	11.30	11.20	11.10	11.00
41	15	11.40	11.30	11.20	11.20	11.00	10.50	10.40	10.20
42	16	11.30	11.20	11.10	11.00	10.40	10.30	10.10	9.50
43	17	11.20	11.10	11.00	10.40	10.20	10.00	9.30	9.00
44	18	11.10	11.00	10.40	10.20	10.00	9.40	9.00	8.10
45	19	11.00	10.40	10.20	10.00	9.40	9.10	8.20	7.30
46	20	10.50	10.30	10.10	9.40	9.20	8.50	7.50	6.50
47	21	10.40	10.20	10.00	9.30	9.00	8.30	7.30	6.10
48	22	10.30	10.10	9.50	9.20	8.40	8.10	7.00	5.40
49	23	10.20	10.10	9.40	9.10	8.30	8.00	6.50	5.10
50	24	10.20	10.00	9.40	9.00	8.20	7.50	6.30	4.50
51	25	10.20	10.00	9.40	9.00	8.20	7.40	6.30	4.40
52	26	10.20	10.00	9.40	9.00	8.20	7.50	6.30	4.40



## Hygiene

- Set up the farm at a safe distance from other poultry houses and fence in.
- Keep birds of only one age group on the farm.
- Keep no other poultry on the farm.
- Allow no visitors to enter the farm.
- Wear only the farm's own protective clothing within the farm area.
- Provide the farm's own protective clothing for veterinarians, service and maintenance workers and consultants.
- Disinfect boots before entering the houses.
- Use bulk feed if possible. Do not allow the truck driver to enter the houses.
- Safeguard the houses against wild birds and vermin. Keep rats and mice under constant control.
- Dispose of dead birds hygienically. Follow local laws and regulations.

## Insect and Parasite Control

If necessary, use a suitable insecticide immediately after the birds have been

removed in order to kill the insects before they are hiding in walls and parts of the equipment.

Use a contact insecticide before warming up the house to control remaining insects. Make sure that the application is safe for the chicks!

## Cleaning and Disinfection

- Disinfection is only effective after cleaning thoroughly.
- Remove litter first and wash floor, walls and all equipment with plenty of water.
- Remove all feed from silos and wash silos. Clean feeding and drinker systems.
- Disinfect walls and floor first time by spraying an effective disinfectant.
- Disinfect smaller parts of the equipment by dipping into a disinfecting solution.
- Disinfect feeders and drinkers by spraying or rinsing with a disinfectant.
- Disinfect the silos by spraying a disinfectant.
- Clean and disinfect streets on the farm area.
- Leave the house for a rest period of 1 – 2 weeks.



## GENERAL RECOMMENDATIONS



- Disinfect house and equipment for a second time by fumigating or fogging a disinfectant.
- Rinse drinker lines with plenty of water before placing chicks.

### Daily Control

Check at least once daily:

- Health status
- Temperature
- Ventilation
- Feed and water consumption
- Lighting
- Mortality

When assessing the state of health, do not just go by the general impression and mortality rate, but also take note of feed and water consumption as well as the consistency of droppings.

### Mating Ratio

Depending on environmental conditions, the mating ratio ranges between 8 and 10 males per 100 females. In controlled environment, usually 8 – 9 males per 100 females are sufficient. In open houses, on slatted floors and hot climate 9 – 10 males per 100 females are rec-

ommended. More males than needed cause disturbance in the flock, resulting in reduced fertility.

### Water Supply

Clean water is as equally important as good feed for top performance. Therefore fresh, clean, potable water must be available at all times for the layers and an adequate consumption must always be assured.

A water meter is a very useful tool to monitor water consumption. The optimal water temperature is about 20 °C (68 °F). Furthermore feed and water intake are closely correlated.

If birds don't drink enough water for any reason, feed intake is consequently reduced. The water to feed ratio at comfortable temperature is around 1.8 : 1, but this relation increases up to 5 : 1 at high ambient temperatures above 30 °C (86 °F). During exposure to high temperatures, birds consume less feed, but more water in an effort to cool their body down.

Check the water quality regularly, especially if you use your own water supply like well water. For example excessive salt levels in drinking water can cause persistent damage to shell quality and hard water with high TDS\* levels may cause kidney damage.

\* TDS: Total Dissolved Solids

## GENERAL RECOMMENDATIONS



### Grit

Feeding grit is not a must but is recommended when rations are supplemented by grains. This stimulates the development of the crop and the gizzard during the rearing period, which in turn has a positive effect on feed intake capacity.

**Table 16: Amount and Granulation of Grit Dependent on Age**

Week 1 – 2	once a week 1 g / bird (size 1 – 2 mm / 0.04 – 0.08 in)
Week 3 – 8	once a week 2 g / bird (size 3 – 4 mm / 0.12 – 0.16 in)
From week 9	once a month 3 g / bird (size 4 – 6 mm / 0.16 – 0.24 in)

### Stocking Density

The optimal bird density depends on management conditions and to which extent climate can be controlled. 6 – 8 birds/m<sup>2</sup> can be taken as a general guide for barn systems. Follow the local laws and statutory regulations.

### Equipment Requirements

In general, the more closely the growing house and facilities resemble the future production system, the easier it will be for the pullets to settle down in their new environment after transfer to the laying house. The following tables show the equipment requirements for rearing and production period.

**Table 17: Equipment Requirement for Rearing Period**

Equipment	Age in Weeks	Requirement
Chick founts	1	1 fount (4 – 5 l) for 100 chicks
Round drinkers	to 20	1 drinker (Ø 46 cm) for 125 birds
Linear drinkers	to 20	1 running m for 100 birds
Nipple drinkers	to 20	6 – 8 birds per nipple
Chick feeding trays	1 – 2	1 tray for 60 chicks
Cut off chick cartons	1 – 2	1 carton for 100 chicks
Round feeders	3 – 10 11 – 20	2 feeders (Ø 40 cm) for 100 birds 3 feeders (Ø 40 cm) for 100 birds
Chain feeders	3 – 10 11 – 20	2.5 – 3.5 lin. m for 100 birds 4.5 lin. m for 100 birds

■ Follow the Instructions of Manufacturer!

## GENERAL RECOMMENDATIONS



Table 18: Equipment Requirement for Production Period

Equipment	Requirement
Round drinkers	1 drinker (Ø 46 cm) for 125 birds
Linear drinkers	1 running m for 80 – 100 birds
Nipple drinkers	4 – 6 birds per nipple
Round feeders	4 feeders (Ø 40 cm) for 100 birds
Single nests	1 nest (26 x 30 cm) for 4 birds
Trough Chain	5 lin. m for 100 birds
Family type nests*	1 sqm for 120 – 130 females

■ Follow the Instructions of Manufacturer!

### Litter

Softwood shavings or straw make suitable litter. Use shavings from untreated wood only in order to avoid poisonings and residues in the egg. Provide sufficient ventilation to ensure good litter condition and remove wet litter, if necessary.

### Nests and Egg Collection

Quality of nests is also a factor which affects egg quality. Regularly renew the litter in litter-type nests and keep them clean. Provide individual nests at a rate of one nest for 4 hens. Collect floor eggs frequently to keep their rate as low as possible. In addition to sufficient nesting space in family type nests, the following

factors are important for a low rate of floor eggs:

- frequent collection of floor eggs
- no draught in the nest area
- clean, dry litter or soft nest lining
- only one type of nest in the house
- easy access
- even distribution of the nests within the house

For optimum hatching egg quality, rollaway nests in combination with slats are better than litter-type nests or family type nests.

## GENERAL RECOMMENDATIONS



### Hatching Egg Care

- Collect hatching eggs frequently.
- Keep floor eggs separately – many will already be contaminated internally.
- If floor eggs are to be used, set in separate incubators (large hatchery) or set at the bottom of setter and hatch trolleys where exploding eggs will cause less damage.
- Take out heavily soiled eggs, do not send them to the hatchery.
- Do not wash hatching eggs.
- Store eggs in a clean egg store. If farm store does not have a controlled temperature, transfer as soon as possible to the hatchery.
- Store eggs at 22 °C (71.6 °F) if setting within 4 days, or 16– 18 °C (60.8 – 64.4 °F) if storing for 5 – 12 days. Older eggs will have markedly lower hatchability.

### Hatching Egg Disinfection

- Fog or spray eggs with a modern disinfectant after collection, then place in store. Follow manufacturers' instructions carefully.
- If desired, fog eggs in egg store once a day, but this should not be necessary if the store is regularly cleaned.
- Eggs should be fogged again before pre-warming and setting.
- Several manufacturers are producing modern disinfectants suitable for use in hatcheries.

- A fogging machine is a good investment as there is no wetting of the eggs and the fog will reach all the eggs.
- Spraying can be carried out using a small droplet size but the spray will not reach all eggs without a fan to aid circulation.
- Formaldehyde (Formalin) is no longer recommended as it is harmful to the embryo, increasing early embryonic death, and it is hazardous to human health.

If fumigation is the only way to disinfect eggs, it can be done in two ways:

- Mixing of 21 cm<sup>3</sup> of Formaldehyde with 17 g of Potassium Permanganate and 21 cm<sup>3</sup> of water per m<sup>3</sup> air
  - **Caution: Never add Potassium Permanganate to Formaldehyde!**
- Heat in of 8 g Paraformaldehyde plus 20 cm<sup>3</sup> of water per m<sup>3</sup> in an electrical heater. Follow closely the instructions of the manufacturer.
  - **Never fumigate with Formaldehyde within the first 96 hours of incubation!**
  - **Never exceed a fumigation time of 30 minutes!**
  - **Keep in mind that the ventilation of a fumigation chamber has to be done with clean air to avoid a recontamination of the hatching eggs!**

## GENERAL INFORMATION



Table 19: Body Weight Development of **LOHMANN LSL** Parent Stock

Age in Weeks	Females			Males
	Average (g)	Range (g)		Average (g)
1	65	62	68	70
2	130	123	137	140
3	190	180	200	200
4	250	237	263	260
5	320	303	337	340
6	400	379	421	430
7	485	460	510	530
8	570	540	600	660
9	650	616	684	785
10	730	692	768	905
11	810	767	853	1020
12	880	834	926	1120
13	945	895	995	1210
14	1005	952	1058	1295
15	1065	1009	1121	1375
16	1120	1061	1179	1450
17	1170	1109	1231	1520
18	1220	1156	1284	1585
19	1270	1203	1337	1645
20	1320	1251	1389	1720
21	1370	1298	1442	1790
22	1420	1345	1495	1855
23	1460	1383	1537	1915
24	1500	1421	1579	1970
25	1535	1454	1616	2020
26	1565	1483	1647	2065
27	1590	1507	1673	2105
28	1610	1525	1695	2140
29	1630	1544	1716	2170
30	1640	1554	1726	2195
31	1648	1561	1735	2215
32	1654	1567	1741	2225
33	1658	1571	1745	2230
34	1660	1573	1747	2234
35	1662	1575	1749	2237
36	1664	1577	1751	2240

## GENERAL INFORMATION



Table 20: Body Weight Development of **LOHMANN BROWN** Parent Stock

Age in Weeks	Females			Males
	Average (g)	Range (g)		Average (g)
1	65	62	68	65
2	130	124	137	145
3	180	171	189	245
4	250	238	263	345
5	320	304	336	445
6	410	390	431	555
7	500	475	525	675
8	590	561	620	795
9	680	646	714	915
10	770	732	809	1035
11	860	817	903	1155
12	950	903	998	1275
13	1030	979	1082	1395
14	1110	1055	1166	1515
15	1190	1131	1250	1635
16	1270	1207	1334	1755
17	1350	1283	1418	1870
18	1440	1368	1512	1985
19	1530	1454	1607	2100
20	1600	1520	1680	2210
21	1650	1568	1733	2290
22	1695	1610	1780	2360
23	1735	1648	1822	2420
24	1773	1684	1862	2475
25	1808	1718	1898	2525
26	1828	1737	1919	2570
27	1843	1751	1935	2610
28	1855	1762	1948	2645
29	1862	1769	1955	2675
30	1870	1777	1964	2695
31	1873	1779	1967	2710
32	1876	1782	1970	2724
33	1879	1785	1973	2739
34	1882	1788	1976	2753
35	1885	1791	1979	2768
36	1888	1794	1982	2782

## GENERAL INFORMATION



Table 21: Performance Goals of **LOHMANN LSL** Parent Stock

Week 21 – 46

Age in Weeks	Rate of Lay (%)		Egg No.		Hatching Eggs			Hatch (%)		No. Saleable Chicks	
	per H.H.	per H.D.	per Weekk	accumulated	%	per Week	accumulated	total Chicks	saleable Chicks	per Week	accumulated
21	30.0	30.0	2.1	2.1	0	0.0	0.0	0	0.0	0.0	0.0
22	55.0	55.0	3.9	6.0	30	1.2	1.2	50	25.0	0.3	0.3
23	73.0	73.1	5.1	11.1	60	3.1	4.2	75	37.5	1.1	1.4
24	83.0	83.2	5.8	16.9	70	4.1	8.3	77	38.5	1.6	3.0
25	88.0	88.3	6.2	23.0	80	4.9	13.2	79	39.5	1.9	5.0
26	90.0	90.4	6.3	29.3	85	5.4	18.6	81	40.5	2.2	7.1
27	91.0	91.5	6.4	35.7	90	5.7	24.3	83	41.5	2.4	9.5
28	91.2	91.8	6.4	42.1	93	5.9	30.2	84	42.0	2.5	12.0
29	91.3	91.9	6.4	48.5	94	6.0	36.2	85	42.5	2.6	14.5
30	91.4	92.1	6.4	54.9	95	6.1	42.3	86	43.0	2.6	17.2
31	91.4	92.2	6.4	61.3	96	6.1	48.5	87	43.5	2.7	19.8
32	91.4	92.3	6.4	67.7	96	6.1	54.6	87	43.5	2.7	22.5
33	91.3	92.3	6.4	74.1	96	6.1	60.7	88	44.0	2.7	25.2
34	91.2	92.3	6.4	80.4	96	6.1	66.9	88	44.0	2.7	27.9
35	91.0	92.2	6.4	86.8	96	6.1	73.0	87	43.5	2.7	30.6
36	90.9	92.2	6.4	93.2	96	6.1	79.1	87	43.5	2.7	33.2
37	90.7	92.2	6.3	99.5	96	6.1	85.2	87	43.5	2.7	35.9
38	90.5	92.1	6.3	105.8	96	6.1	91.3	86	43.0	2.6	38.5
39	90.3	92.0	6.3	112.2	96	6.1	97.3	86	43.0	2.6	41.1
40	90.1	92.0	6.3	118.5	96	6.1	103.4	86	43.0	2.6	43.7
41	89.9	91.9	6.3	124.8	96	6.0	109.4	86	43.0	2.6	46.3
42	89.6	91.8	6.3	131.0	96	6.0	115.4	85	42.5	2.6	48.8
43	89.4	91.6	6.3	137.3	95	5.9	121.4	85	42.5	2.5	51.4
44	89.1	91.5	6.2	143.5	95	5.9	127.3	85	42.5	2.5	53.9
45	88.8	91.3	6.2	149.7	95	5.9	133.2	84	42.0	2.5	56.4
46	88.4	91.1	6.2	155.9	95	5.9	139.1	84	42.0	2.5	58.8

## GENERAL INFORMATION



Table 21: Performance Goals of **LOHMANN LSL** Parent Stock

Week 47 – 72

Age in Weeks	Rate of Lay (%)		Egg No.		Hatching Eggs			Hatch (%)		No. Saleable Chicks	
	per H.H.	per H.D.	per Weekk	accumulated	%	per Week	accumulated	total Chicks	saleable Chicks	per Week	accumulated
47	88.1	91.0	6.2	162.1	95	5.9	144.9	83	41.5	2.4	61.3
48	87.8	90.8	6.1	168.2	95	5.8	150.8	83	41.5	2.4	63.7
49	87.4	90.6	6.1	174.3	94	5.7	156.5	82	41.0	2.4	66.0
50	87.0	90.4	6.1	180.4	94	5.7	162.2	82	41.0	2.3	68.4
51	86.5	90.1	6.1	186.5	94	5.7	167.9	82	41.0	2.3	70.7
52	86.1	89.8	6.0	192.5	94	5.7	173.6	81	40.5	2.3	73.0
53	85.6	89.5	6.0	198.5	93	5.6	179.2	81	40.5	2.3	75.3
54	85.1	89.2	6.0	204.5	93	5.5	184.7	81	40.5	2.2	77.5
55	84.5	88.8	5.9	210.4	93	5.5	190.2	80	40.0	2.2	79.7
56	84.0	88.4	5.9	216.2	92	5.4	195.6	80	40.0	2.2	81.9
57	83.4	88.0	5.8	222.1	92	5.4	201.0	80	40.0	2.1	84.0
58	82.8	87.6	5.8	227.9	92	5.3	206.3	79	39.5	2.1	86.1
59	82.1	87.2	5.7	233.6	92	5.3	211.6	79	39.5	2.1	88.2
60	81.4	86.6	5.7	239.3	92	5.2	216.8	78	39.0	2.0	90.3
61	80.7	86.1	5.6	245.0	92	5.2	222.0	78	39.0	2.0	92.3
62	79.9	85.4	5.6	250.6	92	5.1	227.2	77	38.5	2.0	94.3
63	79.0	84.8	5.5	256.1	92	5.1	232.3	77	38.5	2.0	96.2
64	78.1	84.0	5.5	261.5	91	5.0	237.2	76	38.0	1.9	98.1
65	77.2	83.2	5.4	266.9	91	4.9	242.2	76	38.0	1.9	100.0
66	76.2	82.4	5.3	272.3	90	4.8	247.0	75	37.5	1.8	101.8
67	75.1	81.5	5.3	277.5	90	4.7	251.7	75	37.5	1.8	103.6
68	74.0	80.6	5.2	282.7	89	4.6	256.3	74	37.0	1.7	105.3
69	72.9	79.6	5.1	287.8	89	4.5	260.8	73	36.5	1.7	106.9
70	71.7	78.6	5.0	292.8	88	4.4	265.2	72	36.0	1.6	108.5
71	70.4	77.4	4.9	297.8	88	4.3	269.6	71	35.5	1.5	110.0
72	69.1	76.3	4.8	302.6	87	4.2	273.8	70	35.0	1.5	111.5

## GENERAL INFORMATION



Table 22: Performance Goals of **LOHMANN BROWN** Parent Stock

Week 21 – 46

Age in Weeks	Rate of Lay (%)		Egg No.		Hatching Eggs			Hatch (%)		No. Saleable Chicks	
	per H.H.	per H.D.	per Weekk	accumulated	%	per Week	accumulated	total Chicks	saleable Chicks	per Week	accumulated
21	20.0	20.0	1.4	1.4	0	0.0	0.0	0	0	0.0	0.0
22	35.0	35.1	2.5	3.9	50	1.2	1.2	70	34	0.4	0.4
23	60.0	60.2	4.2	8.1	60	2.5	3.7	75	37	0.9	1.3
24	80.0	80.3	5.6	13.7	70	3.9	7.7	78	38	1.5	2.8
25	88.0	88.4	6.2	19.8	80	4.9	12.6	80	39	1.9	4.8
26	90.0	90.5	6.3	26.1	85	5.4	17.9	82	40	2.1	6.9
27	91.0	91.6	6.4	32.5	90	5.7	23.7	83	40	2.3	9.2
28	91.5	92.2	6.4	38.9	93	6.0	29.6	86	42	2.5	11.7
29	91.7	92.5	6.4	45.3	94	6.0	35.7	88	43	2.6	14.3
30	91.7	92.6	6.4	51.7	95	6.1	41.8	88	43	2.6	16.9
31	91.7	92.8	6.4	58.1	95	6.1	47.9	88	43	2.6	19.5
32	91.6	92.8	6.4	64.6	95	6.1	54.0	88	43	2.6	22.1
33	91.5	92.8	6.4	71.0	95	6.1	60.0	88	43	2.6	24.8
34	91.3	92.7	6.4	77.3	95	6.1	66.1	88	43	2.6	27.4
35	91.1	92.7	6.4	83.7	96	6.1	72.2	88	43	2.6	30.0
36	90.8	92.6	6.4	90.1	96	6.1	78.3	88	43	2.6	32.6
37	90.6	92.4	6.3	96.4	96	6.1	84.4	88	43	2.6	35.2
38	90.3	92.3	6.3	102.7	96	6.1	90.5	88	43	2.6	37.8
39	90.0	92.1	6.3	109.0	96	6.0	96.5	88	43	2.6	40.4
40	89.6	91.9	6.3	115.3	96	6.0	102.5	88	43	2.6	43.0
41	89.3	91.7	6.2	121.5	95	5.9	108.5	88	43	2.5	45.5
42	88.9	91.5	6.2	127.8	95	5.9	114.4	88	43	2.5	48.1
43	88.5	91.3	6.2	134.0	95	5.9	120.3	88	43	2.5	50.6
44	88.0	91.0	6.2	140.1	95	5.9	126.1	88	43	2.5	53.1
45	87.6	90.7	6.1	146.2	95	5.8	131.9	87	42	2.5	55.6
46	87.1	90.4	6.1	152.3	95	5.8	137.7	87	42	2.5	58.0

## GENERAL INFORMATION



Table 22: Performance Goals of **LOHMANN BROWN** Parent Stock

Week 47 – 72

Age in Weeks	Rate of Lay (%)		Egg No.		Hatching Eggs			Hatch (%)		No. Saleable Chicks	
	per H.H.	per H.D.	per Weekk	accumulated	%	per Week	accumulated	total Chicks	saleable Chicks	per Week	accumulated
47	86.6	90.1	6.1	158.4	95	5.8	143.5	86	42	2.4	60.4
48	86.0	89.7	6.0	164.4	95	5.7	149.2	86	42	2.4	62.8
49	85.5	89.4	6.0	170.4	95	5.7	154.9	85	41	2.4	65.2
50	84.9	89.0	5.9	176.3	95	5.6	160.5	85	41	2.3	67.5
51	84.3	88.6	5.9	182.2	94	5.5	166.1	84	41	2.3	69.8
52	83.6	88.1	5.9	188.1	94	5.5	171.6	83	40	2.2	72.0
53	83.0	87.7	5.8	193.9	94	5.5	177.0	82	40	2.2	74.2
54	82.3	87.2	5.8	199.6	94	5.4	182.4	80	39	2.1	76.3
55	81.6	86.7	5.7	205.4	94	5.4	187.8	80	39	2.1	78.4
56	80.8	86.1	5.7	211.0	94	5.3	193.1	80	39	2.1	80.5
57	80.1	85.5	5.6	216.6	94	5.3	198.4	80	39	2.1	82.5
58	79.3	84.9	5.5	222.2	93	5.2	203.6	80	39	2.0	84.5
59	78.5	84.3	5.5	227.7	93	5.1	208.7	80	39	2.0	86.5
60	77.6	83.6	5.4	233.1	92	5.0	213.7	80	39	1.9	88.5
61	76.8	82.9	5.4	238.5	92	4.9	218.6	80	39	1.9	90.4
62	75.9	82.1	5.3	243.8	92	4.9	223.5	78	38	1.9	92.3
63	75.0	81.4	5.2	249.0	92	4.8	228.3	78	38	1.8	94.1
64	74.0	80.6	5.2	254.2	92	4.8	233.1	78	38	1.8	95.9
65	73.1	79.7	5.1	259.3	90	4.6	237.7	76	37	1.7	97.6
66	72.1	78.9	5.0	264.4	90	4.5	242.2	76	37	1.7	99.3
67	71.1	78.0	5.0	269.3	90	4.5	246.7	76	37	1.7	101.0
68	70.0	77.1	4.9	274.2	90	4.4	251.1	74	36	1.6	102.6
69	69.0	76.2	4.8	279.1	90	4.3	255.4	74	36	1.6	104.1
70	67.9	75.3	4.7	283.8	90	4.3	259.7	72	35	1.5	105.6
71	66.7	74.2	4.7	288.5	90	4.2	263.9	72	35	1.5	107.1
72	65.5	73.1	4.6	293.1	90	4.1	268.1	70	34	1.4	108.5



## NOTES

How LOHMANN TIERZUCHT is calculating the energy content of feed and raw materials (International WPSA-formula):

$$\begin{aligned} \text{ME MJ/kg} = & \quad \text{g crude protein} \times 0.01551 \\ & + \text{g crude fat} \times 0.03431 \\ & + \text{g crude starch} \times 0.01669 \\ & + \text{g sugar} \times 0.01301 \text{ (as Saccharose)} \end{aligned}$$

**ME = metabolizable energy in MJ/kg**  
**1 kcal = 4.187 kJ**

Consultation and diagnostics in all questions of poultry health through:

### The LOHMANN TIERZUCHT Veterinary-Laboratory

- Diagnostics
- Quality Control
- Research and Development



**LOHMANN  
TIERZUCHT**

#### **Veterinary-Laboratory**

Abschnede 64 · D-27472 Cuxhaven · Germany  
Phone: +49 (0) 4721 707-244 · Fax: +49 (0) 4721 707-267  
Email: [vetlab@ltz.de](mailto:vetlab@ltz.de) · [www.ltz.de](http://www.ltz.de)

### **Disclaimer**

The information, advices and suggestions given in this management guide should be used for guidance and educational purposes only, recognizing that local environmental and disease conditions may vary and a guide cannot cover all possible circumstances. While every attempt has been made to ensure that the information presented is accurate and reliable at the time of publication, Lohmann Tierzucht cannot accept responsibility for any errors, omissions or inaccuracies in such information or management suggestions.

Further, Lohmann Tierzucht does not warrant or make any representations or guarantees regarding the use, validity, accuracy, or reliability of, or flock performance or productivity resulting from the use of, or otherwise respecting, such information or management suggestions. In no event Lohmann Tierzucht be liable for any special, indirect or consequential damages or special damages whatsoever arising out of or in connection with the use of the information or management suggestions containing in this management guide.



**LOHMANN**  
**TIERZUCHT**

LOHMANN TIERZUCHT GmbH  
Am Seedeich 9-11 • 27472 Cuxhaven • Germany  
P.O. Box 460 • 27454 Cuxhaven • Germany  
Phone +49 (0) 4721 505-0 • Telefax +49 (0) 4721 388-52  
Email: [info@ltz.de](mailto:info@ltz.de) • [www.ltz.de](http://www.ltz.de)